



The effect of short-term meteorological disturbances on the submerged aquatic vegetation and associated fauna in the Patos Lagoon estuary, southern Brazil: a dataset.

Recebido: 16/07/2021 | Aceito: 21/10/2021 | Publicado: 12/11/2021
<https://doi.org/10.53805/lads.v1i2.26>

Dairana Misturini^{*1,2}, Marianna Lanari³, Valéria Lemos⁴, Leonir André Colling⁵

ABSTRACT

Occluded fronts are naturally cyclogenic areas that have been intensified by global warming. Studies evaluating the effects of occluded fronts in the submerged aquatic vegetation (SAV) and its associated fauna in shallow estuarine areas may provide insights on the impacts of climate change-induced extreme weather events on coastal ecosystems functioning. The present dataset describes data on benthic fauna and flora in the Patos Lagoon Estuary (PLE), in southern Brazil, seasonally obtained during intense occluded fronts. Using a hierarchical sample design, based on Beyond BACI protocols (Before/After and Control/Impact), fauna and flora were sampled before and after four occluded fronts passage throughout 2019. Three habitats were sampled: SAV Meadow, SAV Edge and adjacent Sandflat. A total of 432 macrozoobenthic samples (216 samples for stratum); 216 samples for bellow and aboveground biomass, vegetation coverage, canopy height and marine macrophytes morphology; 144 samples by sedimentology and organic matter; 72 water column depth measurements; and 8 temperature and salinity measurements were collected during the study period. The data is available at the Global Biodiversity Information Facility (GBIF), in Darwin Core standard format (DwC), organized according to the OBIS-ENV-DATA model, with CC-BY-NC-4-0 license for use. The present dataset adds to the comprehension of the temporal variability of estuarine benthic communities in subtropical systems, and how short-term meteorological process can affect zoo and fitobenthic communities in the context of climate changes.

Keywords: Benthic Fauna; Seagrass; Sandflat; Occluded Front; Sedimentary Organic Matter.

¹ Programa de Pós-graduação em Oceanografia Biológica – Universidade Federal do Rio Grande, Rio Grande, RS, Brasil. dairana.dai@gmail.com.

² Universidade Federal de Santa Catarina, Coordenadoria Especial de Oceanografia, Laboratório de Biodiversidade Costeira, Florianópolis, SC, Brasil.

³ Universidade Federal do Rio Grande, Laboratório de Ecologia Vegetal Costeira, Rio Grande, Brasil.

⁴ Universidade Federal do Rio Grande, Laboratório de Ictiologia, Rio Grande, Brasil.

⁵ Universidade Federal de Rio Grande, Laboratório de Ecologia de Macroinvertebrados Bentônicos, Rio Grande, Brasil.

PRIOR PUBLICATIONS

MISTURINI, D. A Influência de Sistemas Frontais sobre as Assembleias Bentônicas de Pradarias de Fanerógamas Submersas Estuarinas (Master Dissertation, Universidade Federal do Rio Grande). 2021.

MISTURINI, DAIRANA; COLLING, L. A. Can short-term meteorological events alter subtropical estuarine macrobenthic assemblages in seagrass meadows (Patos Lagoon Estuary-Southern Brazil)?. *Estuarine, Coastal and Shelf Science*, p. 107532, 2021. DOI: <https://doi.org/10.1016/j.ecss.2021.107532>

DATA IMPORTANCE

- The dataset provides information about temporal variation in subtropical estuarine habitats;
- Integration of these data with other similar datasets can add to the comprehension of the effects of SAV on benthic macrofaunal diversity and abundance in subtropical estuarine areas;
- The data have information about different quali and quantitative variations generated by four occluded fronts, in different scenarios of intensity and habitat complexity, for macrozoobenthic organisms and canopy characteristics;
- The dataset offer insights on the influence of patchiness, composition, morphology of seagrass species and edge effect in the structure of macrozoobenthic assemblage.

MATERIALS AND METHODS

Study area

The Patos Lagoon is the largest coastal lagoon in South America with $\sim 11,000$ km². The estuarine area (~ 170 km²; hereafter PLE) comprising shallow flats (> 1.5 m depth) and a deeper main channel (CALLIARI; CUNHA; ANTIQUEIRA, 2010). Located between the latitudes 31° and 32°S, the Patos Lagoon estuary (PLE) (Fig. 1) is situated in a humid subtropical region, with the frequency of occluded fronts passages being variable between seasons (STECH; LORENZZETTI, 1992). The hydrodynamics and physicochemical characteristics of PLE are highly dependent of local and remote winds action, fluvial discharge (mainly by rivers Guaíba, Camaquã and São Gonçalo channel) and regional precipitation (MOLLER et al., 2001). The estuarine region is dominated by riverine discharges and becomes oligohaline in raining season (winter and spring). During Summer and Autumn, local hydrology is mainly controlled by NE and SW wind and eurihaline scenarios prevail (MOLLER et al., 2001).

Field sampling


This study is a short-term research demanded by the Brazilian Long-Term Ecological Research Program (BR-LTER) in the Patos Lagoon Estuary and adjacent marine coast (PLEA). The study was performed in the northern margin of the Justino Bay (Fig. 1, detail), a shallow estuarine area with local hydrodynamics and physical-chemical parameters strongly affects by southwestern winds. The sampled area was surrounded by salt marshes, sandflats and, similarly to other estuarine shoals (SEELIGER; ODEBRECHT, 2010), also presented seasonal variations in the SAV structure and abundance.

The field works were carried out hours or one day before and after four occluded fronts passage, using before as control, according to BACI model (Before/After, Control/Impact; UNDERWOOD, 1996). Sampling fieldwork were in Summer (B: 2019/03/08 and A: 2019/03/13), Autumn (B: 2019/05/08 and A: 2019/05/13), Winter (B: 2019/08/23 and A: 2019/08/28) and Spring (B: 2019/11/08 and A: 2019/11/18). Sampling followed a hierarchical model where submerged aquatic vegetation (SAV) Meadow, Sandflat and

SAV Edge habitats were sorted. The sampled habitats were divided in transects (T1, T2, T3),

creating a 400 m² (20 m x 20 m) delimited squares.

Figure 1. South America, Brazil and Rio Grande do Sul (top left), where the Patos Lagoon Estuary is located (PLE; top left) and the sampling area in the Justino Bay– SJ (shaded in black).



Biological Parameters

During each sampling event, SAV visual coverage and canopy height (quadrat, 1 m²), and biomass (core: 0.1 m Ø X 0.1 m depth) were collected in each transect and habitat (N total = 216 samples). In the laboratory, plants fresh biomass was separated from macrobenthic organisms and sorted out in *R. maritima*/ *Z. palustris* and in *P. striatus* according to leaf morphology. Below (roots and rhizomes) and aboveground (shoots and leaves) biomass were separated for dry weight determination (48 h, 60 °C; MCKENZIE et al., 2003).

Benthic macrofauna were sampled using a cylindrical core (0.008 m²). The sedimentary package was removed from the core, placed in a tray and stratified in the 0.1 m surface sample and 0.1 m to 0.2 m bottom sample, with a spatula, for each transect and habitat (N total = 216 samples for each stratum). The samples were sieved through a 300 µm mesh and kept fresh just to pre-sort fauna and flora. In the laboratory, the samples were identified and classified to the

lowest possible taxonomic level based on morphological and anatomical characteristics, was fixed with 4 % formalin, identified and preserved in alcohol 70 %.

Environmental Parameters

Surface water temperature (mercury thermometer) and salinity (portable RH0-90 refractometer) were taken at the beginning of each field sampling. Sediment samplings were performed in duplicate for each transects and habitat with a cylindrical core (0.002 m², 0.1 m depth, total N=144) for granulometric analyses and determination of organic matter content. Granulometry was analyzed using the method described by Suguio (1973). Organic matter content was determined by the loss on ignition method (DAVIES, 1974).

Quality control

The sampled material was processed by specialists, based on accepted and applied

manuals for sampling (LANA et al., 2006; TURRA; DENADAI, 2015), taxonomy (RIOS, 1994; ARENZON, 1999; AMARAL; RIZZO; ARRUDA, 2006; BUCKUP; BOND-BUCKUP; LARKUM; ORTH; DUARTE, 2006) and methodology (SUGUIO, 1973; DAVIES, 1974; LAVERY; KENDRICK, 2001). Taxonomic validity was verified using the World Register of Marine Species (WoRMS; www.marinespecies.org) and in National Center for Biotechnology Information (<https://www.ncbi.nlm.nih.gov>).


DATA DESCRIPTION

This dataset contains environmental data, morphology and characteristics of SAV data, and benthic macrofauna quali and quantitative data. A total of 432 macrozoobenthic samples (216

samples for stratum); 216 samples for biomass, vegetal coverage, canopy height and marine macrophytes morphology; 144 samples by sedimentology and organic matter; 72 water column depth measurements; and 8 temperature and salinity measurements were seasonally collected. After meteorological events the data differed from the initial (i.e., Before treatments), in this way, we suggest the use of data before events for studies aiming to compare estuarine benthic communities structure and composition.

VAS showed a seasonal trend with higher values in Summer (EV1) and lowers in Winter (EV3). The presence of *R. maritima* was registered throughout the year and *P. striatus* only in Summer (EV1) and Spring (EV4). All vegetation data were expressed per meter or square meter.

Figure 2. Frequency of macrozoobenthic organisms classes in this dataset. "Other" refers to classes whose total density was less than 625 indiv.m⁻².



Our equipment aimed capture benthic macrofauna and marine macrophytes, but we registered other SAV associated fauna occurrence. The associated fauna was constituted by abundant meiofauna, megafauna, limnic organisms, terrestrial insects, planktonic organisms, the complete taxonomic coverage are

in Table 1. A total of 13.828 brackish water macrozoobenthic organisms, distributed across 20 taxa. A total 890 epifaunal organisms and 12.938 infaunal organisms were collected and the most frequent class were the Polychaeta (91 %) and Gastropoda (5 %; Fig 2). The number of organisms is expressed in density (m²).

Table 1. List of *taxa* found in the sampling area during 2019. Note that *taxa* were identified at distinct taxonomic levels. The first column represents the species identifiers numbers at the end of the occurrenceID code in dataset. Sp. ID: Specie Identification in occurrenceID; K: Kingdom; P: Phylum; C: Class; O: Order; F: Family; G: Genus; S: Specie.

Sp ID	K	P	C	O	F	G	S	
3	Animalia	Annelida	Polychaeta	Phyllodocida	Nephtyidae	Nephtys	<i>Nephtys fluviatilis</i>	
4					Nereididae	Alitta	<i>Alitta succinea</i>	
1						Laeonereis	<i>Laeonereis acuta</i>	
16				Spionida	Spionidae	Paraprionospio	<i>Paraprionospio pinnata</i>	
14				Capitellida	Capitellidae	Capitella	<i>Capitella natoi</i>	
2						Heteromastus	<i>Heteromastus similis</i>	
15				Oligochaeta				
38		Arthropoda	Insecta		Coleoptera			
11					Diptera	Chironomidae		
35					Hemiptera	Cicadellidae		
39					Hemiptera			
25					Hymenoptera			
36					Odonata			
41					Malacostraca	Cumacea	Diastylidae	Diastylis
42			Decapoda	Portunidae		Callinectes		
9				Varunidae		Cyrtograpsus	<i>Cyrtograpsus angulatus</i>	
34				Penaeidae				
26			Isopoda	Sphaeromatidae		Cassinidea	Cassinidea	<i>fluminensis</i>
18				Munnidae		Uromunna	<i>Uromunna peterseni</i>	
7			Mysidacea					
8			Amphipoda					
28			Tanaidacea	Kalliapseudidae		Monokalliapseudes	<i>Monokalliapseudes Schubarti</i>	
33				Tanaididae		Sinelobus	<i>Sinelobus Stanfordi</i>	
40			Arachnida	Trombidiformes				
17			Hexanauplia	Sessilia	Balanidae			
37								
32			Entognatha					
20			Branchiopoda	Diplostraca				
27			Ostracoda					
43			Chordata	Thaliacea	Salpida	Salpidae		
19		Cnidaria	Hydrozoa					
5		Mollusca	Gastropoda	Littorinimorpha	Cochliopidae	Heleobia	<i>Heleobia australis</i>	
13							<i>Heleobia charruana</i>	
6								
23	Bivalvia		Myida	Corbulidae	Erodona	<i>Erodona mactroides</i>		
22			Cardiida	Solecurtidae	Tagelus	<i>Tagelus plebeius</i>		
21								
31	Nematoda							
24	Nemertea							
29	Chromista	Foraminifera	Tubothalamea	Miliolida				
30								
46	Plantae	Tracheophyta	Magnoliopsida	Alismatales	Potamogetonaceae	Potamogeton	<i>Potamogeton striatus</i>	
44				Ruppiceae	Ruppia	<i>Ruppia maritima</i>		

Sp ID	K	P	C	O	F	G	S
45					Zannichelliaceae	Zannichellia	<i>Zannichellia palustris</i>

Dataset

The dataset was formatted according to the Darwin Core standard (DwC) and organized in the OBIS-ENV-DATA model (DE POOTER et al., 2017). The file contain an event and location spreadsheet (Event core; Table 2), a species occurrence list spreadsheet (Occurrence extension) and a quantitative environmental

parameters, benthic density and SAV biomass, coverage, morphology and canopy height data spreadsheet (ExtMoF). Darwin Core terms used for each field column is listed in Table 2. Environmental parameters were described with Natural Environment Research Council (NERC).

Table 2. DwC terms and their meanings according to the Darwin Core vocabulary.

Spreadsheet	Field	Definition
Event core	datasetName	Dataset unique identifier
	institutionID	The name of institution having custody of the dataset.
	institutionCode	The name of institution having custody of the dataset.
	rightsHolder	An organization managing the data rights.
	parentEventID	Identifier for Event that groups the Events.
	eventID	A unique identifier constructed based on event information.
	eventDate	The date of field work resented in the Year-Month format.
	Locality	The locality in Patos Lagoon which sampling events occurred.
	locationID	An identifier for locality and habitat which field work are made.
	Country	The country in which sampling events occurred.
	countryCode	The standard code for country.
	stateProvince	The state in which sampling events occurred.
	waterBody	The name of the water body where sampling occurred.
	Habitat	The habitat which sampling events occurred.
	samplingProtocol	Type of sampler method, which defines the sampling protocol.
	samplingEffort	The measure for effort that was expended during a sampling event.
	sampleSizeValue	The size of the sample.
	sampleSizeUnit	Sampler size transformation (0.008 m ²) to obtain organisms density.
geodeticDatum	Spatial reference system upon which the geographic coordinates.	
decimalLatitude	The geographic latitude of sampling area.	
decimalLongitude	The geographic longitude of the sampling area.	
Occurrence extension	basisOfRecord	The origin of the data, such as museum or sample.
	occurrenceID	A unique identifier providing information of biological occurrence.
	taxonRank	The most specific name in the scientific name field.
	occurrenceStatus	A statement about the presence or absence of a Taxon.
	scientificName	The full scientific name, with authorship and date
	scientificNameID	Number identifier of organism
	scientificNameAuthorship	The authorship information for the scientificName.
Kingdom	The kingdom in which the taxon is classified.	

Spreadsheet	Field	Definition
	Phylum	The phylum or division in which the taxon is classified.
	Class	The class in which the taxon is classified.
	Order	The order in which the taxon is classified.
	Family	The family in which the taxon is classified.
	Genus	The genus in which the taxon is classified.
	specificEpithet	The species in which the taxon is classified.
	collectionCode	The laboratory where the collection was deposited
	identifiedBy	Name of the author who identified the organisms.
	recordedBy	The responsible for recording the original Occurrence.
ExtMoF	bibliographicCitation	The bibliographic reference must be cited when the data is used.
	measurementID	Identifier related to the type of measurement performed.
	measurementType	The measurement name type according to NERC.
	measurementTypeID	The identifier for measurement type according to BODC.
	measurementValue	The value of the measurement.
	measurementvalueID	The measurement value ID.
	measurementUnit	The units related to measurement value according to NERC.
	measurementUnitID	A measurement unit according to the BODC
	measurementMethod	The method used to determine the measurement.
	Remarks	Reference or protocol used to obtain the measurements.

NERC: Natural Environment Research Council; BODC: British Oceanographic Data Center

The abbreviations created to identify place, occurrence and measurement in dataset, such as are in the diagram below (Fig. 3).


SUPPLEMENTARY MATERIALS

Repository: Global Biodiversity Information Facility (GBIF). | DOI: <https://doi.org/10.15468/v2dd3g>
 Links to access: <https://www.gbif.org/dataset/04167aa4-4148-4691-a264-031922765fc1>

ACKNOWLEDGEMENTS

The Brazilian Long-Term Ecological Research Program has been supported by MCTI (Ministério de Ciência, Tecnologia e Inovação) and the Brazilian governmental funding agencies CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico, Process NUMBER: CNPq/CAPES/FAPs/BC – Fundo Newton/PELD nº 15/2016), CAPES (Coordenação de Aperfeiçoamento Pessoal de Nível Superior), FAPERGS (Fundação de Amparo à Pesquisa do Rio Grande do Sul), and the Newton Funding (British Council), the Organization for the Conservation of South American Aquatic Mammals. Additional support came from CNPq, which provided research fellowships to author D.M. V.M.L was financially supported by CAPES with a postdoctoral scholarship.

Figure 3. Schema with abbreviations to identify dataset IDs. Codes for strata (10, 20), replicates (a, b, c), transects (t1, t2, t3), habitats (sf, e, sm), periods before (B1, B2, B3, B4) and after (A1, A2, A3, A4) events (2019:3, 2019:5, 2019:8, 2019:11), sampled at Justino Bay (SJ), PLE (patos), BR, as part of the LTER-PLEA and deposited at Benthic Ecology Laboratory (benthos), Oceanographic Institute (IO), Federal University of Rio Grande (FURG). Highlight for the biotic, in green background (top), and abiotic data, in orange (bottom). The red asterisks mark each measurementID specification.



REFERENCES

- AMARAL, A. C. Z.; RIZZO, A. E.; ARRUDA, E. P. Manual de identificação dos invertebrados marinhos da região sudeste-sul do Brasil. [s.l.] EdUSP, v. 1, 2006.
- BUCKUP, L.; BOND-BUCKUP, G.; ARENZON, A. Os crustáceos do Rio Grande do Sul. Porto Alegre: Editora da Universidade, Universidade Federal do Rio Grande do Sul, 1999.
- CALLIARI, L.; CUNHA, R. P. DA; ANTIQUEIRA, J. A. F. DE. Geomorfologia e Dinâmica Sedimentar. In: SEELIGER, U.; ODEBRECHT, C. (Eds.). . O estuário da Lagoa dos Patos: um século de transformações. 1. ed. Rio Grande: FURG, p. 31–43. 2010.
- DAVIES, B. E. Loss-on-ignition as an estimate of soil organic matter. Soil Science Society of America Journal, v. 38, n. 1, p. 150–151, jan. 1974.
- DE POOTER, D. et al. Toward a new data standard for combined marine biological and environmental datasets - Expanding OBIS beyond species occurrences. Biodiversity Data Journal, v. 5, n. 1, 2017.
- LANA, P. DA C. et al. Avaliação ambiental de estuários brasileiros: diretrizes metodológicas. serie livr ed. Rio de Janeiro: Museu Nacional, 2006.
- LARKUM, A. W. D.; ORTH, R. J.; DUARTE, C. Seagrasses: Biology, Ecology and Conservation. 1. ed. Dordrecht: Springer Netherlands, 2006.

LAVERY, P.; KENDRICK, G. Assessing biomass, assemblage structure and productivity of algal epiphytes on seagrasses. 2001.

MCKENZIE, L. J.; CAMPBELL, S. J.; RODER, C. A. Seagrass-Watch: manual for mapping & monitoring seagrass resources by community (citizen) volunteers. 2. ed. Queensland: Seagrass-Watch: Manual for Mapping & Monitoring Seagrass Resources by Community (citizen) volunteers, 2003.

MOLLER, O. O. et al. The influence of local and non-local forcing effects on the subtidal circulation of Patos Lagoon. *Estuaries*, v. 24, n. 2, p. 297, 2001. DOI: <https://doi.org/10.2307/1352953>.

RIOS, E. C. Seashells of Brazil. 2. ed. Rio Grande: Fundação Cidade do Rio Grande, Fundação Universidade do Rio Grande, Museu Oceanográfico, 1994.

SEELIGER, U.; ODEBRECHT, C. O Estuário da Lagoa dos Patos: um século de transformações. 1. ed. Rio Grande: Universidade Federal do Rio Grande - FURG, 2010.

STECH, J. L.; LORENZZETTI, J. A. The Response of the South of Wintertime Brazil Bight to the Passage Cold Fronts. *Journal Of Geophysical Research*, v. 97, n. 6, p. 9507–9520, 1992. DOI: <https://doi.org/10.1029/92JC00486>.

SUGUIO, K. Introdução à sedimentologia. São Paulo: E. Blücher, 1973.

TURRA, A.; DENADAI, M. R. Protocolos para o monitoramento de habitats bentônicos costeiros - Rede de Monitoramento de Habitat Bentônicos Costeiros - ReBentos. [s.l.] Instituto Oceanográfico da Universidade de São Paulo, 2015.

UNDERWOOD, A. J. On Beyond Baci. In: SCHMITT, R. J.; OSENBURG, C. W. B. T.-D. E. I. (Eds.). *Detecting Ecological Impacts*. San Diego: Elsevier, 1996. p. 151–175.